

CSF biomarkers of Alzheimer disease in HIV-associated neurologic disease



D.B. Clifford, MD
A.M. Fagan, PhD
D.M. Holtzman, MD
J.C. Morris, MD
M. Teshome, MD
A.R. Shah, MS
J.S.K. Kauwe, PhD

Address correspondence and reprint requests to Dr. David B. Clifford, Department of Neurology, Washington University in St. Louis, Box 8111, 660 South Euclid, St. Louis, MO 63110
cliffordd@neuro.wustl.edu

ABSTRACT

Background: HIV-associated neurologic disorders (HAND) continue to develop in many patients with HIV. CSF amyloid measurements in HAND have been reported to be similar to those in dementia of the Alzheimer type (DAT). Confirmatory evaluation of this finding in carefully evaluated subjects is needed.

Methods: CSF specimens were obtained from subjects clinically categorized with normal cognition from the general population, HIV+ subjects with normal cognition, HIV+ subjects with impaired cognition, or presumed HIV– subjects with mild DAT. CSF measurements of β -amyloid_(1–42) (A β 42), β -amyloid_(1–40) (A β 40), total tau (t-tau), and phosphorylated tau (p-tau181) were performed.

Results: CSF A β 42 measured in 49 HAND subjects had a median level of 501 pg/mL, which was lower than that of 50 controls of similar age who had median of 686 pg/mL ($p < 0.0001$) or 21 HIV+ subjects without cognitive impairment who had median of 716 pg/mL ($p < 0.003$). HAND subjects had similar CSF A β 42 to 68 subjects with mild DAT. There was no difference of CSF A β 40 between the groups. Tau and p-tau181 was elevated in DAT, but slightly lower than control in both HIV+ groups.

Conclusions: β -Amyloid_(1–42) (A β 42) measurements in CSF of cognitively impaired patients with HIV are similar to those in patients with mild dementia of the Alzheimer type (DAT). Normal or slightly depressed CSF tau and p-tau181 measurements distinguish these patients with HIV-associated neurologic disorders (HAND) from patients with DAT. Further evaluation of amyloid metabolism in patients with HIV cognitive disorder is needed to understand the implications of depressed CSF A β 42 in the setting of HAND. *Neurology*® 2009;73:1982–1987

GLOSSARY

A β = β -amyloid; AD = Alzheimer disease; APP = amyloid precursor protein; CDR = Clinical Dementia Rating; CHARTER = CNS Highly Activated Retroviral Therapy Effects Research; DAT = dementia of the Alzheimer type; HAD = HIV-associated dementia; HAND = HIV-associated neurologic disorders; LRP = lipoprotein receptor related protein; MCD = mild cognitive disorder; NNTC = National NeuroAIDS Tissue Consortium; p-tau181 = phosphorylated tau; t-tau = total tau.

HIV-associated neurocognitive disorders (HAND) continue to be a problem in HIV-infected (HIV+) subjects despite generally successful virologic control with modern antiretroviral agents. The pathophysiology of observed deficits in HAND is incompletely understood, but appears to be enhanced by aging, consistent with an interaction of aging seen for other neurodegenerative processes.¹ Measurements of CSF β -amyloid_(1–42) (A β 42), total tau (t-tau), and phosphorylated tau (p-tau181) in advanced HAND, Alzheimer disease (AD), and age-matched seronegative controls have been reported previously.² HAND subjects had significantly decreased CSF A β 42 and increased t-tau and p-tau181 concentrations similar to patients with AD, suggesting that HAND may be associated with AD or an AD-like process. Additional observations linking HAND and amyloid-associated processes include reports of neuropatho-

Editorial, page 1942

See also page 1975

e-Pub ahead of print on November 11, 2009, at www.neurology.org.

From the Department of Neurology (D.B.C., A.M.F., D.M.H., J.C.M., M.T., A.R.S.), Washington University School of Medicine, St. Louis, MO; and Department of Biology (J.S.K.K.), Brigham Young University, Provo, UT.

Supported by NS32228 and NIMH 22005 (D.B.C.), P50 AG05681, P01 AG026276, and P01 AG03991 (J.C.M., D.M.H., A.M.F.), NNTC: NS38841 (Singer), R24MH59724 (Morgello), R24NS45491 (Gelman), MH05745 (Grant).

Disclosure: Author disclosures are provided at the end of the article.

logic evidence of amyloid deposition in the brains of HIV+ patients,³ of more silver stained (i.e., neuritic) amyloid plaques in HIV+ patients than in controls of the same age,⁴ and a demonstration of increased diffuse plaques in HIV brains.⁵ It has been demonstrated that in neural cell membranes from human brain aggregates, the HIV-1 transactivator regulatory protein Tat may inhibit neprilysin, an endonuclease for A β breakdown, the primary peptide constituent of amyloid in AD.⁵

Given the expanding insights for therapeutics to address abnormal A β metabolism in AD, clear demonstration of the role of similar pathophysiologic properties in HAND could be of substantial importance. Previous reports² are provocative, but incomplete since this article did not report A β 40, which specifically remains normal when A β 42 declines in AD.⁶ Specific reductions in CSF A β 42 have been shown to be associated with brain amyloid deposition as evidenced by PET imaging of the amyloid-binding agent Pittsburgh Compound B.⁷ The interpretation of decreased levels of CSF A β 42 as being due to possible AD processes would be different in the face of concomitant decreases in A β 40. Further, the findings of the previous report² regarding tau levels in HAND are at variance to several prior studies.^{8,9} In the present study, we compared CSF values of A β and tau within long-term infected HIV+ subjects with known cognitive status to age-matched HIV uninfected controls and patients diagnosed with mild dementia of the Alzheimer type (DAT).

METHODS Subjects. CSF from clinically characterized HIV+ patients was obtained from both the National NeuroAIDS Tissue Consortium (NNTC) and from the Washington University in St. Louis cohort of the CNS Highly Activated Retroviral Therapy Effects Research (CHARTER) study. All subjects gave written informed consent under the direction of the local institutional review boards for human studies. For both cohort studies from which data were used, all subjects had neuropsychometric and neurologic examinations leading to classification of cognitive impairment. NNTC subjects were enrolled due to advanced HIV disease with anticipated imminent death. All 41 CSF samples from the NNTC cohort had HAND, either classified as HIV-associated dementia (HAD) (n = 11) or mild cognitive disorder (MCD) (n = 30), both of which require impaired performance on a broad neuropsychometric battery and evidence of functional impairment. Within the CHARTER co-

hort, an additional 8 subjects had HAND, with the rest (n = 21) having normal performance on concurrent neurologic examination and neuropsychometric testing. Diagnostic classification of HAND is not made in the face of serious comorbid conditions that might be alternative explanations for neurocognitive performance deficits.

Controls were derived from a clinically well-characterized group of adults volunteering for evaluation at the Washington University Memory and Aging Project. Careful examination and testing was performed with participants categorized by the Clinical Dementia Rating (CDR) scale as unimpaired (CDR = 0) or having mild impairment that has a high correlation with histologic AD in subsequent neuropathologic examinations (CDR = 0.5).⁶

CSF handling and evaluation. CSF samples obtained from the NNTC had been rapidly frozen and stored and shipped frozen to Washington University for analysis. The control samples were placed on ice immediately during the lumbar puncture, and stored frozen until time of analysis. HIV samples were also quickly frozen and thawed only once for the analysis. Control and DAT samples were collected at 7:30 AM, while the times of collection for the HIV samples were random daytime hours.

CSF samples were analyzed for total tau, phospho-tau₁₈₁ (p-tau181), and A β ₄₂ by commercial ELISA (Innotest, Innogenetics, Ghent, Belgium), and A β ₄₀ by ELISA as previously described by Fagan et al.⁷ For all measures, samples were continuously kept on ice, and assays were performed on sample aliquots after a single thaw following initial freezing.

Statistical analysis. Standard descriptive statistics were calculated for each population in the study. CSF A β 42 and A β 40 measurements were log transformed to approximate a normal distribution. CSF tau and p-tau181 required log-log transformation to approximate a normal distribution. Contrasts of biomarker measurements among the control, DAT, HIV, and HAND groups were performed using analysis of covariance after adjusting for age. *p* Values for pairwise comparisons were taken from the least squares means difference matrix with a significant difference if *p* < 0.05. Correlations among CD4, nadir CD4, plasma viral load, and CSF viral load were calculated using Pearson correlation coefficient. All statistical analyses were performed using SAS version 9.1.

RESULTS Table 1 provides demographic variables for neurologically normal HIV-positive patients, HAND subjects, DAT subjects, and nondemented controls. Overall, the DAT cohort was older than any other group (for all contrasts *p* < 0.0001). We dealt with this difference in age by including age as a covariate in the analyses. Most patients with HIV, but not all, were taking antiretroviral therapy. Within the HIV+ subgroups, HAND patients had a trend toward lower nadir and current CD4 counts and higher plasma viral loads than cognitively normal HIV+ patients. However, none of the differences is significant. Median CSF viral load was higher in the neurologically normal HIV+ subjects. CSF biomarkers results are presented in table 2 and demonstrated in the figure. CSF A β 42 in the HAND HIV group was significantly lower than unaffected controls or neurologically normal HIV+

Table 1 Characteristics of each sample

	No.	Age, y, mean (SD)	CD4, cells/mm ³ , mean (median)	Nadir CD4, cells/mm ³ , mean (median)	Plasma HIV viral load, c/mL, mean (median)	CSF HIV viral load, c/mL, mean (median)
Control	50	50 (3.02)	NA	NA	NA	NA
DAT	68	74 (7.39)	NA	NA	NA	NA
Neuro normal HIV+	21	43 (9.18)	421 (389)	218 (225)	19,317 (13,000)	5,750 (671)
HAND HIV+	49	48 (8.03)	348 (320)	176 (117)	127,067 (452)	4,450 (<50)

Control without dementia (Control), mild (Clinical Dementia Rating 0.5/1.0) dementia of the Alzheimer type (DAT), neurocognitively normal HIV-positive (HIV), and neurocognitively impaired HIV-associated neurologic disorder (HAND) samples are shown.

subjects. Even after statistical corrections for the differences in age between DAT and HAND subjects, the CSF A β 42 levels in HAND subjects are not significantly different from CSF A β 42 levels in DAT subjects ($p = 0.25$). CSF A β 40 levels were similar for all groups. In contrast, CSF tau and p-tau181 (the primary constituents of neurofibrillary tangles in AD) for both neurologically normal and HAND HIV+ groups were slightly lower than unaffected controls, and did not mirror the significant elevations typically seen with DAT (figure).

We also sought to identify possible associations of the biomarker findings with important measures of the biology of HIV disease including current CD4 count, nadir CD4 count, current plasma viral load, and current CSF viral load. Pearson correlation coefficients were derived and revealed no significant correlations of any single HIV disease markers to any of the 4 AD biomarkers we studied (table 3).

DISCUSSION In clinically well-characterized subjects, we find that HIV-positive subjects with neurocognitive impairment have significantly lower CSF A β 42 than unimpaired subjects, comparable to findings in mild DAT. However, we find that the tau elevations characteristic of DAT are absent in HAND. Our findings suggest that there is a change in A β 42 metabolism occurring with the development of HIV-associated neurologic dysfunction.


Age-related changes in A β 42 do not account for differences since the HAND and control populations had similar ages, and all analyses include age as a covariate. The observed reduction in CSF A β 42 in cognitively impaired HIV-positive patients is consistent with an earlier report² and, in the absence of reductions in CSF A β 40, suggests that the reductions in A β 42 may be indicative of brain amyloid deposition and not just death of neurons. However, in contrast to an earlier report,² and consistent with other studies,^{8,9} we observed that CSF tau and p-tau181 were not elevated in association with reductions in CSF A β 42. This CSF profile helps distinguish HIV from DAT, implicating abnormalities in amyloid metabolism in HAND, with no evidence of the tauopathy that occurs in DAT. Another recent report¹⁰ shows similar findings with regard to A β 42, associating lower CSF A β 42 but normal p-tau181 with symptomatic HIV-associated cognitive disease. However, this report continues to suggest total tau was elevated in CSF of this population. The reason for the disparity in tau measurements is unclear, although one possible difference is that these patients were untreated for HIV, whereas our symptomatic patients were generally taking antiretroviral therapy. It is possible that participants in this study may have had more active or ongoing neuropathology due to lack of HIV treatment. However, this report of normal tau in the DAT patients is also discordant with AD literature. Together, our findings suggest that measurements of certain tau species in CSF may be useful as a means of differentiating HIV-associated cognitive decline from AD. Because of the aging of the HIV-positive population, this distinction may well be of increasing utility.

Further consideration of the reasons for low CSF A β 42 in HAND require ongoing research. Reductions in CSF A β 42 levels have been shown to be a reliable marker for brain amyloid deposition in patients with AD as well as a subset of cognitively normal elderly individuals hypothesized to have “preclinical” AD (i.e., amyloid deposition prior to

Table 2 Mean (median) values of biomarkers in each sample

	CSF biomarkers			p-tau 181, pg/mL
	A β 42, pg/mL	A β 40, pg/mL	Tau, pg/mL	
Control	722 (686)	10,773 (9,875)	242 (212)	47 (44)
DAT	421 (359)	10,492 (10,128)	572 (475)	85 (71)
Neuro normal HIV	683 (716)	8,267 (7,921)	183 (191)	33 (32)
HAND HIV	522 (501)	9,852 (9,051)	196 (174)	40 (39)

A β 42, A β 40, tau, and p-tau mean and median values for controls without dementia (Control), mild (Clinical Dementia Rating 0.5/1.0) dementia of the Alzheimer type (DAT), neurocognitively normal HIV-positive (HIV), and neurocognitively impaired HIV-associated neurologic disorder (HAND). See figure for significance of contrasts.



(A) Aβ42 levels in HIV-negative controls without dementia (controls), individuals with dementia of the Alzheimer type (DAT), HIV-positive individuals without dementia (HIV), and individuals with HIV-associated neurologic disorders (HAND). *p* Values of less than 0.05 from the least squares means difference matrix are shown. (B) Aβ40 levels in the 4 groups. There are no significant differences. (C) Tau levels in the 4 groups. DAT is significantly higher than all other groups. Both HIV and HAND are significantly lower than controls. No other significant associations. (D) p-tau181 levels in the 4 groups. DAT is significantly higher than controls, HIV, and HAND. The HAND group is significantly lower than controls. No other significant associations.

the onset of clinical dementia symptoms).¹¹⁻¹⁴ While the pathology of HAND and AD differs significantly,¹⁵ it will be important to see if PET amyloid imaging in vivo can distinguish these forms of dementia despite the similarity of the CSF Aβ42 levels. We assume that in HIV, if amyloid deposition is occurring, there will be a much greater percentage of diffuse, nonfibrillar Aβ deposits as compared to what is typically seen in AD, given the absence of florid amyloid plaques in HIV neuropathology.

There are reasons that could explain altered Aβ metabolism in HIV disease. HIV Tat protein has been shown to inhibit an Aβ cleaving protein, neprilysin, which could lead to increased brain amyloid deposition.⁵ Alternatively, HIV Tat may compete with the amyloid precursor protein (APP) and apolipoprotein E (an Aβ chaperone) for binding to the low density lipoprotein receptor related protein (LRP), thus blocking LRP-mediated clearance of Aβ from brain interstitial fluid to peripheral compartments.¹⁶ On the other hand, soluble APP cleavage products (sAPPα and sAPPβ) have been reported to

be reduced in the CSF of patients with HAND compared to those with DAT or HIV-negative controls, with sAPPα showing a slight decline in the asymptomatic HIV state.¹⁷ These findings, in conjunction with the low Aβ42 in HIV, might suggest that synthesis of the amyloid precursor (APP) declines with symptomatic disease. If APP synthesis were decreased, one would expect a decrease in CSF Aβ40 and Aβ42, not just a selective decrease in Aβ42. However, this idea could be tested since Aβ turnover is quite a dynamic process, approximating 8% turnover per hour,¹⁸ and could be assessed directly.

Our findings reinforce the importance of understanding the significant change in amyloid metabolism that is associated with symptomatic HIV-associated neurologic disease and may expand considerations of the pathophysiology of HAND. The failure of HIV viral load and CD4 counts to correlate with development of HAND in our HIV-positive patients reinforces the need for reliable predictive biomarkers for this widespread problem. While CSF viral load was previously re-

Table 3 Pearson correlation coefficients comparing CD4, nadir CD4, plasma HIV viral load, and CSF HIV viral load with the 4 CSF biomarkers				
	CD4	Nadir CD4	Plasma viral load	CSF viral load
Aβ42	0.04967	−0.01721	−0.04829	0.03366
p Value	0.6875	0.8909	0.698	0.7789
No.	68	66	67	72
Aβ40	−0.20002	−0.0902	0.04676	0.19117
p Value	0.1046	0.4749	0.7093	0.1184
No.	67	65	66	68
Tau	−0.17622	−0.20969	−0.02007	−0.00995
p Value	0.1506	0.0911	0.8719	0.9339
No.	68	66	67	72
p-tau181	−0.18255	−0.16487	0.00508	0.00172
p Value	0.1362	0.1859	0.9675	0.9885
No.	68	66	67	72

There are no significant correlations.

lated to development of dementia, in the current treatment era this is no longer the case, as we found in this analysis.^{19,20}

Successful HIV therapy is resulting in long survival of patients with HIV, who are now entering age brackets where AD becomes more common. In aging patients developing dementia, our findings suggest that CSF tau measurements are likely to distinguish patients with HAND from those with AD. While age was included as a covariate in our analyses to account for age-related changes, it may be necessary to confirm that CSF tau distinguishes HAND from DAT once a substantial sample of older patients with HAND comparable in age to DAT samples are available for analysis. Meanwhile, emerging technologies such as assessment of amyloid deposition in the brain with PET imaging and defining the dynamics of A β production and elimination may clarify the basis for the changes being observed.

AUTHOR CONTRIBUTIONS

Statistical analysis was conducted by Dr. John S.K. Kauwe.

ACKNOWLEDGMENT

The authors thank the volunteers providing CSF and time for assessments from the National NeuroAIDS Tissue Consortium, The Memory and Aging Project, and the CHARTER project at Washington University in St. Louis. They also recognize the leadership of the site investigators at NNTC sites including Dr. E. Singer, Dr. S. Morgello, Dr. B. Gelman, and Dr. I. Grant. Michael Spinner performed the A β 40 assay.

DISCLOSURE

The study was funded by NIH grant MH22005. Dr. Clifford serves/has served on scientific advisory boards for Biogen Idec, Elan Corporation, Roche, Forest Laboratories, Inc., Genentech, Inc., GlaxoSmithKline, Millennium Pharmaceuticals, Inc., Schering-Plough Corp., Bristol-Meyers Squibb, and Genzyme Corporation; received speaker honorarium and funding for travel from GlaxoSmithKline; has received research support from Pfizer Inc., Schering-Plough Corp., Bavarian Nordic, NeurogesX, GlaxoSmithKline, Tibotec Therapeutics, Boehringer Ingelheim, and

Gilead Sciences, Inc.; and receives research support from the NIH [U01 NS32228 (PI), U01 AI69495 (PI), NIMH 22005 CHARTER Project (Site PI), NIDA R03 DA022137 (Co-I), NIMH MH058076 (Site PI), and R21 3857-53187 (PI)]. Dr. Fagan serves on a speakers' bureau for the Alzheimer's Association. Dr. Holtzman serves on scientific advisory boards for Satori Pharmaceuticals and EnVivo Pharmaceuticals; serves as an Associate Editor of *Annals of Neurology*, the *Journal of Neuroscience*, *Neurobiology of Disease*, and *Experimental Neurology*; may accrue revenue on pending US Patent 20080145941 (filed 6/18/08): Methods for Measuring the Metabolism of Neurally Derived Biomolecules in Vivo, pending US Patent 20090074775 (filed 3/19/09): Use of Anti-AB Antibody to Treat Traumatic Brain Injury, pending US Patent 20090035298 (filed 2/5/09): Methods to Treat Alzheimer's Disease or Other Amyloid Beta Accumulation Associated disorders; US Patent 7,195,761 (issued 3/27/07): Humanized antibodies that sequester abeta peptide, US Patent 7,015,044 (issued 3/21/06): Diagnostic for early stage Alzheimer's disease, US Patent 6,465,195 (issued 10/15/02): Predictive diagnostic for Alzheimer's disease; serves as a consultant to Merck Serono, Eli Lilly and Company, Takeda Pharmaceutical Company Limited, Abbott, Comentis, Inc., Eisai Inc., and AstraZeneca; is cofounder of and receives board of directors compensation from C2N Diagnostics LLC; receives research support from AstraZeneca, Pfizer Inc., Eli Lilly and Company, Elan Corporation, Forest Laboratories, Inc., the NIH [NIA R37 AG13956 (PI), NINDS 1P30NS057105 (PI), NINDS P01-NS35902 (PI of project 3), NINDS P01-NS32636 (PI of project 3), NIA P01-AG026276 (Co-I), NIA R01-AG025824 (I), NINDS R01-NS034467 (I), NIA U01AG032438 (Co-I), NIA P01-AG03991 (PI of project 2)], Cure Alzheimer's Fund, and Fidelity Foundation; has received compensation from Washington University from license revenue received for licensing of patent application entitled "Methods for Measuring the Metabolism of Neurally Derived Biomolecules in Vivo" to C2N Diagnostics LLC; and may receive future royalty payments for Washington University licensing patent entitled "Methods for Measuring the Metabolism of Neurally Derived Biomolecules in Vivo" to C2N Diagnostics, LLC, and could receive future royalty payments from Washington University for licensing patent entitled "Humanized antibodies that sequester abeta peptide" US Patent 7,195,761 to Eli Lilly and Company. Dr. Morris serves on scientific advisory boards for AstraZeneca, Bristol-Myers Squibb, Genentech, Inc., Merck Serono, Novartis, Pfizer Inc., Schering-Plough Corp., Eli Lilly and Company, Wyeth, and Elan Corporation; serves on the editorial advisory board of *Alzheimer's Disease and Associated Disorders*; receives royalties from publishing *Mild Cognitive Impairment and Early Alzheimer's Disease* (John Wiley and Sons, 2008), *Dementia* (Clinical Publishing, 2007), *Handbook of Dementing Illnesses, 2 Ed* (Taylor & Francis, 2006), and for an editorial in *Lancet Neurology* (Elsevier, 2008); and receives research support from Elan Corporation, Wyeth, Eli Lilly and Company, Novartis, Pfizer Inc., Avid Radiopharmaceuticals, the NIH/NIA [P50AG05681 (PI), P01AG03991 (PI), P01AG026276 (PI), U01AG032438 (PI), U01AG024904 (Neuropathology Core Leader), R01AG16335 (Consultant), and P50NS006833 (Investigator), and from the Dana Foundation. A.R. Shah, Dr. Teshome, and Dr. Kauwe report no disclosures.

Received April 10, 2009. Accepted in final form August 3, 2009.

REFERENCES

1. Valcour V, Shikuma C, Shramizu B, et al. Higher frequency of dementia in older HIV-1 individuals: The Hawaii Aging with HIV-1 cohort. *Neurology* 2004;63: 822–827.
2. Brew BJ, Pemberton L, Blennow K, Wallin A, Hagberg L. CSF amyloid β 42 and tau levels correlate with AIDS dementia complex. *Neurology* 2005;65:1490–1492.
3. Green DA, Masliah E, Vinters HV, Beizai P, Moore DJ, Achim CL. Brain deposition of beta-amyloid is a common pathologic feature in HIV positive patients. *AIDS* 2005; 19:407–411.
4. Esiri MM, Biddolph SC, Morris CS. Prevalence of Alzheimer plaques in AIDS. *J Neurol Neurosurg Psychiatry* 1998;65:29–33.

5. Rempel HC, Pulliam L. HIV-1 Tat inhibits neprilysin and elevates amyloid β . *AIDS* 2005;19:127–135.
6. Fagan AM, Csernansky CA, Morris JC, Holtzman DM. The search for antecedent biomarkers of Alzheimer's disease. *J Alzheimer Dis* 2005;8:347–358.
7. Fagan AM, Mintun MA, Mach RH, et al. Inverse relation between in vivo amyloid imaging load and cerebrospinal fluid $A\beta_{42}$ in humans. *Ann Neurol* 2006;59:512–519.
8. Ellis RJ, Seubert P, Motter R, et al. Cerebrospinal fluid tau protein is not elevated in HIV-associated neurologic disease in humans. *Neurosci Lett* 1998;254:1–4.
9. Green AJE, Giovannoni G, Hall-Craggs MA, Thompson EJ, Miller RF. Cerebrospinal fluid tau concentrations in HIV infected patients with suspected neurological disease. *Sex Transm Inf* 2000;76:443–446.
10. Brew B, Gisslén M, Pemberton L, et al. CSF τ , p - τ and amyloid- β 1–42 in HIV infection. 15th Conference on Retroviruses and Opportunistic Infections 2008;197. Abstract.
11. Rowe CC, Ng S, Ackermann U, et al. Imaging β -amyloid burden in aging and dementia. *Neurology* 2007;68:1718–1725.
12. Fagan AM, Roe CM, Xiong C, Mintun MA, Morris JC, Holtzman DM. Cerebrospinal fluid tau/ β -amyloid $_{42}$ ratio as a prediction of cognitive decline in nondemented older adults. *Arch Neurol* 2007;64:343–349.
13. Mintun MA, LaRossa GN, Sheline YI, et al. [^{11}C]PIB in a nondemented population: potential antecedent marker of Alzheimer disease. *Neurology* 2006;67:446–452.
14. Fagan AM, Head D, Shah AR, et al. Decreased cerebrospinal fluid Abeta (42) correlates with brain atrophy in cognitively normal elderly. *Ann Neurol* 2009;65:176–183.
15. Anthony IC, Ramage SN, Carnie FW, Simmonds P, Bell JE. Accelerated Tau deposition in the brains of individuals infected with human immunodeficiency virus-1 before and after the advent of highly active anti-retroviral therapy. *Acta Neuropathol* 2006;111:529–538.
16. Liu Y, Jones M, Hingtgen CM, et al. Uptake of HIV-1 Tat protein mediated by low-density lipoprotein receptor-related protein disrupts the neuronal metabolic balance of the receptor ligands. *Nat Med* 2000;6:1380–1387.
17. Gisslén M, Blennow K, Brew B, et al. CSF neural marker profile distinguishes AIDS dementia complex from Alzheimer's disease. 15th Conference on Retroviruses and Opportunistic Infections 2008;196. Abstract.
18. Bateman RJ, Munsell LY, Morris JC, Swarm R, Yarasheski KE, Holtzman DM. Human amyloid- β synthesis and clearance rates as measured in cerebrospinal fluid in vivo. *Nat Med* 2006;12:856–861.
19. Cysique LA, Maruff P, Brew BJ. Prevalence and pattern of neuropsychological impairment in human immunodeficiency virus-infected/acquired immunodeficiency syndrome (HIV/AIDS) patients across pre-and post-highly active antiretroviral therapy eras: a combined study of two cohorts. *J Neurovirol* 2004;10:350–357.
20. Robertson KR, Smurzynski M, Parsons TD, et al. The prevalence and incidence of neurocognitive impairment in the HAART era. *AIDS* 2007;21:1915–1921.

Your Voice + 2 Days = The Future of Neurology

Bring your expertise as a neurology professional to Capitol Hill for the Eighth Annual Neurology on the Hill, March 8 and 9, 2010, in Washington, DC. No prior advocacy experience is necessary to apply and on-site training will be provided. The deadline for applications is December 13, 2009. Selected participants will receive updates on current health policy issues and discuss needed improvements in neurologic care with Congressional legislators and staff. For more information or to apply, visit www.aan.com/noh or contact Melissa Larson at mlarson@aan.com or (651) 695-2748.